

Atomic Pair Distribution Function (PDF) and X-Ray Scattering Methods to Assess the Stability of Amorphous Organic Compounds

D. Beckers, M. Gateshki, PANalytical B.V., Almelo, The Netherlands
A. Adibhatla, PANalytical Inc., Westborough MA, USA

The amorphous state is of significant interest as a possible means to enhance aqueous solubility of Active Pharmaceutical Ingredients (APIs). Amorphization is achieved either by spray drying the API itself or by creating solid dispersions of the API with different polymers. Often the short-range structural arrangement directly determines the physical stability of the pharmaceutical solid. An important practical barrier to the development of amorphous APIs for drug products is the lack of reliable methods for structural characterization and fingerprinting. The Atomic Pair Distribution Function (PDF) has been suggested for fingerprinting of amorphous materials and to study the short range order (i.e. inter-atomic distances) of the materials. The PDF technique utilizes a Fourier transformation of the X-ray total scattering data (Bragg peaks and diffuse scattering) and gives information about the inter-atomic distances of the material. The accuracy of this assessment of inter-atomic distances can be improved by using a radiation source producing high-energy X-rays.

Recent advances in laboratory X-ray diffractometer technology like e.g. new generation detectors optimized for hard radiation (like Mo and Ag radiation) as well as improved software algorithms allow to minimize artifacts or fluctuations in the PDF arising from statistical noise, resulting in more reliable data. Thus, the amorphous and nanocrystalline materials in the drugs can be studied reliably in the laboratory. We will demonstrate the possibilities of laboratory PDF on a variety of organic samples of different nature and will show that PDF patterns can reliably be used with cluster analysis for fingerprinting of amorphous drugs and drug compounds.